arXiv:1112.1215v1 [math.CO] 6 Dec 2011

A Note on the Characterization of Digraph Sequences ⋆

Annabell Berger

Department of Computer Science
Martin-Luther-Universität Halle-Wittenberg
`berger@informatik.uni-halle.de`

Abstract. We consider the following fundamental realization problem of directed graphs. Given a sequence $S := \binom{a_1}{b_1}, \ldots, \binom{a_n}{b_n}$ with $a_i, b_i \in \mathbb{Z}_0^+$. Does there exist a digraph (no parallel arcs allowed) $G = (V, A)$ with a labeled vertex set $V := \{v_1, \ldots, v_n\}$ such that for all $v_i \in V$ indegree and outdegree of v_i match exactly the given numbers a_i and b_i, respectively? There exist two known approaches solving this problem in polynomial running time. One first approach of Kleitman and Wang (1973) uses recursive algorithms to construct digraph realizations [KW73]. The second one draws back into the Fifties and Sixties of the last century and gives a complete characterization of digraph sequences (Gale 1957, Fulkerson 1960, Ryser 1957, Chen 1966). That is, one has only to validate a certain number of inequalities. Chen bounded this number by n. His characterization demands the property that S has to be in lexicographical order. We show that this condition is much too strong. Hence, we can give several, different sets of n inequalities. We think that this stronger result can be very important with respect to structural insights about the sets of digraph sequences for example in the context of threshold sequences. The new characterization is fomally analogous to the classical one by Erdős and Gallai (1960) for undirected graphs.

1 Characterization of Digraph Sequences

Problem 11 (digraph realization problem) *Given is a finite sequence* $S := \binom{a_1}{b_1}, \ldots, \binom{a_n}{b_n}$ *with* $a_i, b_i \in \mathbb{Z}_0^+$. *Does there exist a digraph (without parallel arcs)* $G = (V, A)$ *with a labeled vertex set* $V := \{v_1, \ldots, v_n\}$ *such that we have indegree* $d_G^-(v_i) = a_i$ *and outdegree* $d_G^+(v_i) = b_i$ *for all* $v_i \in V$*?*

If the answer is "yes", we call sequence S *digraph sequence* and digraph G a *digraph realization.* We exclude tuples $\binom{0}{0}$ in S. Furthermore, we will tacitly assume that $\sum_{i=1}^n a_i = \sum_{i=1}^n b_i$, as this is obviously a necessary condition for any realization to exist, since the number of ingoing arcs must equal the number of outgoing arcs. We denote sequence S by *decreasing lexicographical sorted* if we have $\binom{a_i}{b_i} \geq_{lex} \binom{a_{i+1}}{b_{i+1}}$ for all $i \in \{1, \ldots, n-1\}$, i.e. $(a_i > a_{i+1}) \vee (a_i = a_{i+1} \wedge b_i \geq b_{i+1})$. There exist two known approaches solving this problem in polynomial running time. One first approach of Kleitman and Wang uses recursive algorithms to construct digraph realizations [KW73]. The second one gives a complete characterization of digraph sequences. Hence, it is possible to check a polynomial number of inequalities (in the size of the number of tuples in a sequence) which leads to the correct decision of the realizability of a sequence. However, these views come from another analogous problem — the *graph realization problem* which asks whether a given *undirected sequence* $S := (d_1), \ldots, (d_n)$ with $d_i \in \mathbb{Z}_0^+$ possesses a realization as a graph. The characterization approach was found by Erdős and Gallai [EG60] and a realization algorithm was introduced by Havel [Hav55] and Hakimi [Hak62]. Four authors, namely David Gale [Gal57], Herbert J. Ryser [Rys57], Delbert Ray Fulkerson [Ful60] and Wai-Kai Chen [Che66] gave sufficient and necessary conditions which completely characterize digraph sequences. Actually, none of the mentioned authors has found the following theorem in its general form. Gale and Ryser dealt with digraph sequences where at most

---

⋆ This work was supported by the DFG Focus Program Algorithm Engineering, grant MU 1482/4-2.

one loop per vertex is allowed. However, their insights are fundamental for characterizing digraph sequences without loops. Fulkerson gave a general result with exponential many inequalities and Chen formulated and proved the final details. Therefore, we cite the following theorem as the result of all four authors.

Theorem 1 (Characterization of digraph sequences [Ful60,Rys57,Che66,Gal57]). *Let* $S := \binom{a_1}{b_1}, \dots, \binom{a_n}{b_n}$ *be a sequence in decreasing lexicographical order.* S *is a digraph sequence if and only if the following conditions are fulfilled:*

$$\sum_{i=1}^{k} \min(b_i, k-1) + \sum_{i=k+1}^{n} \min(b_i, k) \geq \sum_{i=1}^{k} a_i$$

for all $k \in \{1, \dots, n\}$.

In our note, we relax the conditions of this theorem in this sense that we show that sequence S has only to be sorted in decreasing order with respect to its first components a_i. Hence, the condition of a lexicographical sorted sequence is not neccessary and much too strong. We think that this observation could be very important in the context of *threshold sequences* (see the book by Mahadev and Peled[MP95]). A threshold sequence can be defined as this sequence fullfilling all inequalities in Theorem 1 with equality. Hence, our result leads to several and not only one threshold sequence for a given sequence S (dependent on the number of decreasing permutations of S). To prove our claim, we go back to the results of Fulkerson [Ful60], take some ideas of Chen's proof [Che66] and show our more general result. In the following, we denote by S_k a *subsequence of* S consisting of k arbitrary tuples from S and we write $\overline{S_k}$ for the corresponding subsequence containing all remaining tuples of S.

Theorem 2 (Fulkerson 1960). *Sequence* $S := \binom{a_1}{b_1}, \dots, \binom{a_n}{b_n}$ *is a digraph sequence if and only if for all subsequences* S_k *and* $k \in \{1, \dots, n\}$ *we have*

$$\sum_{\{b_i|\ \binom{a_i}{b_i} \text{ tuple in } S_k\}} \min(b_i, k-1) + \sum_{\{b_i|\ \binom{a_i}{b_i} \text{ tuple in } \overline{S_k}\}} \min(b_i, k) \geq \sum_{\{a_i|\ \binom{a_i}{b_i} \text{ tuple in } S_k\}} a_i. \quad (*)$$

Furthermore, we need a simple and nice combinatorial insight.

Proposition 1 (Principle of double counting). *For an abitrary, finite set* $M \subset \mathbb{N}$ *and* $k \in \mathbb{N}$ *we have*

$$\sum_{i \in M} \min(i, k) = \sum_{j=1}^{k} |\{i \in M|\ i \geq j\}|.$$

Proof. Consider Figure 1.

Each $i \in M$ corresponds to a row with i entries "one". Then $\min(i, k)$ counts exactly the number of "ones" in row i which is truncated at length k. Hence, $\sum_{i \in M} \min(i, k)$ sums up all entries "one" in each k-truncated row. On the other hand, we can count these entries by considering all "ones" in columns $t \in \{1, \dots, k\}$. The number of entries in the jth column is exactly the number of rows i with $i \geq j$, i.e., $|\{i \in M|\ i \geq j\}|$. This proves our claim.

□

This insight gives a further useful formulation of Theorem 2.

Corollary 1 (Reformulation of Fulkerson). *Sequence* $S := \binom{a_1}{b_1}, \dots, \binom{a_n}{b_n}$ *is a digraph sequence if and only if for all subsequences* S_k *and* $k \in \{1, \dots, n\}$ *we have*

$$|\{b_i|\ \binom{a_i}{b_i} \text{ tuple in } \overline{S_k}, b_i \geq k\}| + \sum_{j=1}^{k-1} |\{b_i|\ \binom{a_i}{b_i} \text{ tuple in } S, b_i \geq j\}| \geq \sum_{\{a_i|\ \binom{a_i}{b_i} \text{ tuple in } S_k\}} a_i \quad (+)$$



Fig. 1. Rows with i entries "one" (black and blue). $\min(i,k)$ are the black entries in each row corresponding to k columns with entries "one" and "zero".

Proof. Applying the principle of double counting (Proposition 1) for each summand in (*), we get a further formulation for each subsequence S_k with $k \in \{1,\dots,n\}$.

$$\begin{aligned}
\sum_{\{a_i \mid \binom{a_i}{b_i} \text{ tuple in } S_k\}} a_i \quad &\le \sum_{\{b_i \mid \binom{a_i}{b_i} \text{ tuple in } S_k\}} \min(b_i, k-1) + \sum_{\{b_i \mid \binom{a_i}{b_i} \text{ tuple in } \overline{S_k}\}} \min(b_i, k) \\
&\overset{\text{Prop. 1}}{=} \sum_{j=1}^{k-1} \left|\left\{b_i \middle| \begin{pmatrix} a_i \\ b_i \end{pmatrix} \text{ tuple in } S_k, b_i \ge j\right\}\right| + \sum_{j=1}^{k} \left|\left\{b_i \middle| \begin{pmatrix} a_i \\ b_i \end{pmatrix} \text{ tuple in } \overline{S_k}, b_i \ge j\right\}\right| \\
&= \sum_{j=1}^{k-1} \left|\left\{b_i \middle| \begin{pmatrix} a_i \\ b_i \end{pmatrix} \text{ tuple in } S, b_i \ge j\right\}\right| + \left|\left\{b_i \middle| \begin{pmatrix} a_i \\ b_i \end{pmatrix} \in \overline{S_k}, b_i \ge k\right\}\right|
\end{aligned}$$

□

It is easy to derive from Proposition 1 that

$$\sum_{i\in M} \min(i,k) - \sum_{i\in M} \min(i,k+1) \ge 0, \tag{1}$$

because the $(k+1)$th column in our Figure 1 cannot contain more entries then the kth column. We apply this insight for the "left-hand-sides" (*) of the inequalities from Fulkerson's Theorem. For a simpler notion, we denote this left-hand-side of (*) in Fulkerson's Theorem 2 by

$$x(S_k) := \sum_{\{b_i \mid \binom{a_i}{b_i} \text{ tuple in } S_k\}} \min(b_i, k-1) + \sum_{\{b_i \mid \binom{a_i}{b_i} \text{ tuple in } \overline{S_k}\}} \min(b_i, k).$$

Proposition 2. *Let S_{k-1} be an arbitrary subsequence of sequence $S := \binom{a_1}{b_1}, \dots, \binom{a_n}{b_n}$ with $k-1$ tuples, S_k an extension of S_{k-1} containing a further tuple of S and S_{k+1} an extension of S_k by a further tuple of S. Then we have $x(S_{k+1}) - x(S_k) \le x(S_k) - x(S_{k-1}) + 1$.*

Proof.

$$
\begin{aligned}
x(S_{k+1}) - x(S_k) &= \sum_{\{b_i \mid \binom{a_i}{b_i} \text{ tuple in } S_{k+1}\}} \min(b_i, k) + \sum_{\{b_i \mid \binom{a_i}{b_i} \text{ tuple in } \overline{S_{k+1}}\}} \min(b_i, k+1) \\
&\quad - \left(\sum_{\{b_i \mid \binom{a_i}{b_i} \text{ tuple in } S_k\}} \min(b_i, k-1) + \sum_{\{b_i \mid \binom{a_i}{b_i} \text{ tuple in } \overline{S_k}\}} \min(b_i, k) \right) \\
&= \sum_{\{b_i \mid \binom{a_i}{b_i} \text{ tuple in } S_k\}} \min(b_i, k) + \min(b_{k+1}, k) + \sum_{\{b_i \mid \binom{a_i}{b_i} \text{ tuple in } \overline{S_{k+1}}\}} \min(b_i, k+1) \\
&\quad - \left(\sum_{\{b_i \mid \binom{a_i}{b_i} \text{ tuple in } S_k\}} \min(b_i, k-1) + \min(b_{k+1}, k) + \sum_{\{b_i \mid \binom{a_i}{b_i} \text{ tuple in } \overline{S_{k+1}}\}} \min(b_i, k) \right) \\
&= \sum_{\{b_i \mid \binom{a_i}{b_i} \text{ tuple in } S_k\}} \min(b_i, k) - \sum_{\{b_i \mid \binom{a_i}{b_i} \text{ tuple in } S_k\}} \min(b_i, k-1) \\
&\quad + \sum_{\{b_i \mid \binom{a_i}{b_i} \text{ tuple in } \overline{S_{k+1}}\}} \min(b_i, k+1) - \sum_{\{b_i \mid \binom{a_i}{b_i} \text{ tuple in } \overline{S_{k+1}}\}} \min(b_i, k) \\
&\overset{\text{Prop. 1}}{=} \left|\left\{b_i \mid \begin{pmatrix} a_i \\ b_i \end{pmatrix} \text{ tuple in } S_k, b_i \geq k\right\}\right| + \left|\left\{b_i \mid \begin{pmatrix} a_i \\ b_i \end{pmatrix} \text{ tuple in } \overline{S_{k+1}}, b_i \geq k+1\right\}\right| \\
&\leq \left|\left\{b_i \mid \begin{pmatrix} a_i \\ b_i \end{pmatrix} \text{ tuple in } S_k, b_i \geq k-1\right\}\right| + \left|\left\{b_i \mid \begin{pmatrix} a_i \\ b_i \end{pmatrix} \text{ tuple in } \overline{S_{k+1}}, b_i \geq k\right\}\right| \\
&= \left|\left\{b_i \mid \begin{pmatrix} a_i \\ b_i \end{pmatrix} \text{ tuple in } S_{k-1}, b_i \geq k-1\right\}\right| + \left|\{b_k \mid b_k \geq k-1\}\right| \\
&\quad + \left|\left\{b_i \mid \begin{pmatrix} a_i \\ b_i \end{pmatrix} \text{ tuple in } \overline{S_k}, b_i \geq k\right\}\right| - \left|\{b_{k+1} \mid b_{k+1} \geq k\}\right| \\
&\leq \left|\left\{b_i \mid \begin{pmatrix} a_i \\ b_i \end{pmatrix} \text{ tuple in } S_{k-1}, b_i \geq k-1\right\}\right| + 1 + \left|\left\{b_i \mid \begin{pmatrix} a_i \\ b_i \end{pmatrix} \text{ tuple in } \overline{S_k}, b_i \geq k\right\}\right| \\
&\overset{\text{Prop. 1}}{=} \sum_{\{b_i \mid \binom{a_i}{b_i} \text{ tuple in } S_{k-1}\}} \min(b_i, k-1) - \sum_{\{b_i \mid \binom{a_i}{b_i} \text{ tuple in } S_{k-1}\}} \min(b_i, k-2) + 1 \\
&\quad + \left(\sum_{\{b_i \mid \binom{a_i}{b_i} \text{ tuple in } \overline{S_k}\}} \min(b_i, k) - \sum_{\{b_i \mid \binom{a_i}{b_i} \text{ tuple in } \overline{S_k}\}} \min(b_i, k-1) \right) \\
&= \sum_{\{b_i \mid \binom{a_i}{b_i} \text{ tuple in } S_{k-1}\}} \min(b_i, k-1) + \min(b_k, k-1) + \sum_{\{b_i \mid \binom{a_i}{b_i} \text{ tuple in } \overline{S_k}\}} \min(b_i, k) + 1 \\
&\quad - \left(\sum_{\{b_i \mid \binom{a_i}{b_i} \text{ tuple in } S_{k-1}\}} \min(b_i, k-2) + \min(b_k, k-1) + \sum_{\{b_i \mid \binom{a_i}{b_i} \text{ tuple in } \overline{S_k}\}} \min(b_i, k-1) \right) \\
&= \sum_{\{b_i \mid \binom{a_i}{b_i} \text{ tuple in } S_k\}} \min(b_i, k-1) + \sum_{\{b_i \mid \binom{a_i}{b_i} \text{ tuple in } \overline{S_k}\}} \min(b_i, k) + 1 \\
&\quad - \left(\sum_{\{b_i \mid \binom{a_i}{b_i} \text{ tuple in } S_{k-1}\}} \min(b_i, k-2) + \sum_{\{b_i \mid \binom{a_i}{b_i} \text{ tuple in } \overline{S_{k-1}}\}} \min(b_i, k-1) \right) \\
&= x(S_k) - x(S_{k-1}) + 1
\end{aligned}
$$

□

The idea of Chen was to show that it is sufficient to consider only the n subsequences $S_k := \binom{a_1}{b_1}, \ldots, \binom{a_k}{b_k}$ with $k \in \{1, \ldots, n\}$. The only requirement is that sequence S is sorted in decreasing lexicographical order. Since, the proof of Chen omitted the consideration of several case distinctions, we here repeat his idea and extend it to our general scenario in adding several own insights.

Theorem 3 (Digraph Characterization). *Sequence* $S := \binom{a_1}{b_1}, \ldots, \binom{a_n}{b_n}$ *with* $a_1 \geq \cdots \geq a_n$ *is a digraph sequence if and only if we find for all* $k \in \{1, \ldots, n\}$ *that*

$$\sum_{i=1}^{k} \min(b_i, k-1) + \sum_{i=k+1}^{n} \min(b_i, k) \geq \sum_{i=1}^{k} a_i. \tag{**}$$

Proof. $\Rightarrow$: Since S is a digraph sequence, we can apply Theorem 2 of Fulkerson. Condition (*) is fullfilled for all subsequences S_k. Obviously, the n subsequences $S_k := \binom{a_1}{b_1}, \ldots, \binom{a_k}{b_k}$ with $k \in \{1, \ldots, n\}$ are such subsequences leading to our condition (**).
$\Leftarrow$: We have to show that conditions (**) lead to conditions (*) in Theorem 2. In this case, sequence S is a digraph sequence. Clearly, condition (**) fullfills condition (*) for $S_k := \binom{a_1}{b_1}, \ldots, \binom{a_k}{b_k}$ with $k \in \{1, \ldots, n\}$.

We prove the theorem in two steps.

1. We show that it is sufficient to consider in (*) **all** subsequences $S_k^j := \binom{a_{j_1}}{b_{j_1}}, \ldots, \binom{a_{j_k}}{b_{j_k}}$ of sequence S where $a_{j_1} \geq \cdots \geq a_{j_n}$ is fullfilled for S_k^j and its corresponding sequence $\overline{S_k^j}$. Clearly, j ranges between 1 and the number l of different possible sortings of S with respect to the first component a_i. (We avoid isomorphic sortings.) Hence, we get $n \cdot l$ inequalities.
2. We show that it is sufficient to consider in (*) **for one arbitrary** $j \in \{1, \ldots, l\}$ the n subsequences $S_k^j := \binom{a_{j_1}}{b_{j_1}}, \ldots, \binom{a_{j_k}}{b_{j_k}}$ of sequence S where $a_{j_1} \geq \cdots \geq a_{j_n}$ is fullfilled for S_k^j and its corresponding sequence $\overline{S_k^j}$.

Step 1. We consider two permutations of S, namely S^j and $S^{j'}$ which differ in the position $\mu < \kappa$ ($\mu, \kappa \in \{1, \ldots, n\}$) of exactly two different tuples, i.e. $\binom{a_{j_\mu}}{b_{j_\mu}} = \binom{a_{j'_\kappa}}{b_{j'_\kappa}}$ and $\binom{a_{j_\kappa}}{b_{j_\kappa}} = \binom{a_{j'_\mu}}{b_{j'_\mu}}$. S^j shall fullfill $a_{j_1} \geq \cdots \geq a_{j_n}$ and $S^{j'}$ not, because we want to show: If S_k^j fullfills for each k condition (*) in Theorem 2, then $S_k^{j'}$ does it, too. So, we can assume that $a_{j_\mu} > a_{j_\kappa}$. Let us now use the equivalent formulation (+) of Fulkerson (Corollary 1). Note that two such inequalities (+) for one given k – one for sequence S^j and another for $S^{j'}$ – can only differ, when $\mu \leq k < \kappa$. We denote the second summand of (+) by $K(S_k) := \sum_{j=1}^{k-1} |\{b_i|\ \binom{a_i}{b_i} \text{ tuple in } S, b_i \geq j\}|$. Clearly, we have $K(S_k^j) = K(S_k^{j'})$ for each k. We consider the remaining summands and denote them for simplicity by $B(S_k) := \{b_i|\ \binom{a_i}{b_i} \in \overline{S_k}, b_i \geq k\}$ and

$$A(S_k) := \sum_{\{a_i|\ \binom{a_i}{b_i} \text{ tuple in } S_k\}} a_i.$$

With $\mu \leq k < \kappa$ and $a_{j_\mu} > a_{j_\kappa}$ we get $A(S_k^{j'}) \leq A(S_k^j) - 1$. Let us consider all $k \in \{1, \ldots, n\}$ with $b_{j_\mu}, b_{j'_\mu} \geq k$ or $b_{j_\mu}, b_{j'_\mu} < k$, respectively. We either find $b_{j_\kappa} \in B(S_k^j), b_{j'_\kappa} \in B(S_k^{j'})$ or $b_{j_\kappa} \notin B(S_k^j), b_{j'_\kappa} \notin B(S_k^{j'})$, respectively. We get $|B(S_k^{j'})| = |B(S_k^j)|$. Hence, we get by our assumption for this case

$$K(S_k^{j'}) + |B(S_k^{j'})| = K(S_k^j) + |B(S_k^j)| \geq A(S_k^j) > A(S_k^{j'}).$$

So, it remains to distinguish the cases

(1) $b_{j_\mu} \geq k > b_{j'_\mu}$ or
(2) $b_{j'_\mu} \geq k > b_{j_\mu}$.

In case (1) we find $b_{j'_\kappa} \geq k > b_{j_\kappa}$. It follows $b_{j_\kappa} \notin B(S^j_k), b_{j'_\kappa} \in B(S^{j'}_k)$. We get $|B(S^{j'}_k)| = |B(S^j_k)| + 1 \geq |B(S^j_k)|$. Hence, we get

$$K(S^{j'}_k) + |B(S^{j'}_k)| = K(S^j_k) + |B(S^j_k)| + 1 \geq A(S^j_k) + 1 \geq A(S^{j'}_k).$$

In case (2) we get $b_{j_\kappa} \geq k > b_{j'_\kappa}$. It follows $b_{j_\kappa} \in B(S^j_k), b_{j'_\kappa} \notin B(S^{j'}_k)$. This results in $|B(S^{j'}_k)| = |B(S^j_k)| - 1$. We get

$$K(S^{j'}_k) + |B(S^{j'}_k)| = K(S^j_k) + |B(S^j_k)| - 1 \geq A(S^j_k) - 1 \geq A(S^{j'}_k).$$

So, we can conclude that for two given permutations $S^j, S^{j'}$ of S which differ only in two positions and fulfill the conditions of this paragraph, it is sufficient to consider the permutation S^j in the Theorem of Fulkerson 2. Let us now consider a sorting S^d of S with $a_{d_1} \geq \cdots \geq a_{d_n}$ and a permutation S^t where $a_{t_1} \geq \cdots \geq a_{t_n}$ is not fullfilled. Furthermore, we assume for all $k \in \{1, \ldots, n\}$ we have $K(S^d_k) + |B(S^d_k)| \geq A(S^d_k)$. Clearly, it is possible to construct a sequence $S^d := S^1, \ldots, S^\nu := S^t$ of sortings of S such that two adjacent permutations S^j and S^{j+1} in this sequence differ only in two positions $\mu < \kappa$ an we have $a_{j_\mu} > a_{j_\kappa}$ in S^j. Then we can conclude for each $k \in \{1, \ldots, n\}$ step by step with our insights above and starting with $K(S^1_k) + |B(S^1_k)| \geq A(S^1_k)$ that $K(S^2_k) + |B(S^2_k)| \geq A(S^2_k), \ldots, K(S^j_k) + |B(S^j_k)| \geq A(S^j_k), \ldots, K(S^t_k) + |B(S^t_k)| \geq A(S^t_k)$. This completes the proof of our step 1.

Step 2. We consider two permutations of S, namely S^j and $S^{j'}$ with $a_{j_1} \geq \cdots \geq a_{j_n}$ and $a_{j'_1} \geq \cdots \geq a_{j'_n}$. These two permutations shall only differ in two adjacent positions μ and $\mu + 1$ ($\mu \in \{1, \ldots, n-1\}$). Clearly, we have $a_{j_\mu} = a_{j'_\mu}$ and $a_{j_{\mu+1}} = a_{j'_{\mu+1}}$ by our assumptions. Hence, we can assume $b_{j_\mu} \neq b_{j'_\mu}$ and $b_{j_{\mu+1}} \neq b_{j'_{\mu+1}}$, respectively. Otherwise, there is nothing to prove. Again, we use the equivalent formulation $(+)$ of Fulkerson (Corollary 1) and show for each k:

$$K(S^j_k) + |B(S^j_k)| \geq A(S^j_k) \Leftrightarrow K(S^{j'}_k) + |B(S^{j'}_k)| \geq A(S^{j'}_k). \qquad (++)$$

Clearly, this claim is fullfilled for all $k \geq \mu + 1$ and for all $k < \mu$, because all summands are equal. Furthermore, we have $A(S^j_k) = A(S^{j'}_k)$ for all k. Hence, we only have to treat the sets $B(S^j_k)$ and $B(S^{j'}_k)$ in the case $\mu = k$. For $b_{j_\mu}, b_{j'_\mu} \geq \mu$ or $b_{j_\mu}, b_{j'_\mu} < \mu$, respectively, we either find $b_{j_{\mu+1}} \in B(S^j_\mu), b_{j'_{\mu+1}} \in B(S^{j'}_\mu)$ or $b_{j_{\mu+1}} \notin B(S^j_\mu), b_{j'_{\mu+1}} \notin B(S^{j'}_\mu)$. We get $|B(S^{j'}_\mu)| = |B(S^j_\mu)|$. Let us now assume that

$$X(S^j_\mu) = K(S^j_\mu) + |B(S^j_\mu)| \geq A(S^j_\mu)$$

and

$$x(S^{j'}_\mu) = K(S^{j'}_\mu) + |B(S^{j'}_\mu)| < A(S^{j'}_\mu).$$

Since, $K(S^j_k) = K(S^{j'}_k)$ and $A(S^j_k) = A(S^{j'}_k)$ for all k this can only happen when $|B(S^j_\mu)| > |B(S^{j'}_\mu)|$. Since, S^j_μ and $S^{j'}_\mu$ do only differ in their μth tuple, this is only possible for the case $b_{j_\mu} < \mu \leq b_{j'_\mu}$. In particular, we have $|B(S^j_\mu)| = |B(S^{j'}_\mu)| + 1$. Hence, with our insights using our notation from above we have

1. $x(S^j_{\mu-1}) = X(S^{j'}_{\mu-1}) \geq A(S^j_{\mu-1}) = A(S^{j'}_{\mu-1})$.
2. $x(S^{j'}_{\mu}) + 1 = X(S^j_{\mu}) \geq A(S^j_{\mu}) = A(S^{j'}_{\mu})$ and $X(S^{j'}_{\mu}) < A(S^{j'}_{\mu})$.
3. $x(S^j_{\mu+1}) \geq A(S^j_{\mu+1})$ and $X(S^{j'}_{\mu+1}) \geq A(S^{j'}_{\mu+1})$.

Since, $|B(S^j_{\mu})| = |B(S^{j'}_{\mu})| + 1$ and $A(S^j_{\mu}) = A(S^{j'}_{\mu})$ we can conclude $x(S^j_{\mu}) = A(S^j_{\mu})$ in 2. But then it follows with 1. $x(S^j_{\mu}) - x(S^j_{\mu-1}) \leq A(S^j_{\mu}) - A(S^j_{\mu-1}) = a_{j_{\mu}}$. On the other hand, it follows from 3. that $x(S^j_{\mu+1}) - x(S^j_{\mu}) \geq a_{j_{\mu}}$. With 1. and 2. we conclude

$$X(S^{j'}_{\mu}) - X(S^{j'}_{\mu-1}) = X(S^j_{\mu}) - 1 - X(S^j_{\mu-1}) \leq a_{j_{\mu}} - 1.$$

and

$$X(S^{j'}_{\mu+1}) - X(S^{j'}_{\mu}) = X(S^j_{\mu+1}) - (X(S^j_{\mu}) - 1) \geq a_{j_{\mu+1}} + 1 = a_{j_{\mu}} + 1.$$

But then it follows $X(S^{j'}_{\mu+1}) - X(S^{j'}_{\mu}) \geq X(S^{j'}_{\mu}) - X(S^{j'}_{\mu-1}) + 2$. Contradiction to Proposition 2! Hence, our assumption was false and it follows $K(S^{j'}_{\mu}) + |B(S^{j'}_{\mu})| \geq A(S^{j'}_{\mu})$. Carefully looking into our arguments the same is true for the other side of our implication (++) (this is the simpler case). Let us now consider two permutations S^d and S^t of S with $a_{d_1} = a_{t_1} \geq \cdots \geq a_{d_n} = a_{t_n}$. Furthermore, we assume for all $k \in \{1, \ldots, n\}$ that we have $K(S^d_k) + |B(S^d_k)| \geq A(S^d_k)$. Clearly, it is possible to construct a sequence $S^d := S^1, \ldots, S^{\nu} := S^t$ of permutations S^j with $a_{j_1} \geq \cdots \geq a_{j_n}$ such that two adjacent sortings S^j and S^{j+1} in this sequence differ only in two adjacent positions $\mu < \mu + 1$. Then we can conclude for each $k \in \{1, \ldots, n\}$ step by step with our insights above and starting with $K(S^1_k) + |B(S^1_k)| \geq A(S^1_k)$ that $K(S^2_k) + |B(S^2_k)| \geq A(S^2_k), \ldots, K(S^j_k) + |B(S^j_k)| \geq A(S^j_k), \ldots, K(S^t_k) + |B(S^t_k)| \geq A(S^t_k)$. This completes the proof of our step 2. ☐

The idea of Chen was to consider all n inequalities of the form $K(S_k) \geq A(S_k) - |B(S_k)|$ with $k \in \{1, \ldots, n\}$. He proved that the right-hand-side of these inequalities is maximized if S is decreasing lexicographical sorted and S_k consists of the first k tuples of S. Unfortunately, by this approach he overlooked that other decreasing sortings of S with respect to the a_i are also sufficient and the condition of a lexicographical sorting is too strong.

References

Che66. W. Chen, *On the realization of a (p,s)-digraph with prescribed degrees*, Journal of the Franklin Institute **281** (1966), no. 5, 406 – 422.

EG60. P. Erdős and T. Gallai, *Graphs with prescribed degree of vertices (hungarian)*, Mat. Lapok **11** (1960), 264–274.

Ful60. D.R. Fulkerson, *Zero-one matrices with zero trace*, Pacific J. Math. **10** (1960), 831–836.

Gal57. D. Gale, *A theorem on flows in networks*, Pacific J.Math. **7** (1957), 1073–1082.

Hak62. S.L. Hakimi, *On the realizability of a set of integers as degrees of the vertices of a simple graph*, J. SIAM Appl. Math. **10** (1962), 496–506.

Hav55. V. Havel, *A remark on the existence of finite graphs*, Casopis Pest. Math. **80** (1955), 477–480.

KW73. D. J. Kleitman and D. L. Wang, *Algorithms for constructing graphs and digraphs with given valences and factors*, Discrete Mathematics **6** (1973), no. 1, 79 – 88.

MP95. N. V. R. Mahadev and U. N. Peled, *Threshold graphs and related topics*, North-Holland, 1995.

Rys57. H.J. Ryser, *Combinatorial properties of matrices of zeros and ones*, Canad J.Math. **9** (1957), 371–377.